CLEO CONF95-15
EPS0174
July 7, 1995

OBSERVATION OF THE Ξ_c^+ CHARMED BARYON DECAYS TO $\Sigma^+K^-\pi^+$, Σ^+K^{*0}, and $\Lambda K^-\pi^+\pi^+$

M. Bishai,[1] J. Fast,[1] E. Gerndt,[1] J.W. Hinson,[1] T. Miao,[1] D.H. Miller,[1] M. Modesitt,[1] E.I. Shibata,[1] I.P.J. Shipsey,[1] P.N. Wang,[1] L. Gibbons,[2] S.D. Johnson,[2] Y. Kwon,[2] S. Roberts,[2] E.H. Thorndike,[2] T.E. Coan,[3] J. Dominick,[3] V. Fadeyev,[3] I. Korolkov,[3] M. Lambrecht,[3] S. Sanghera,[3] V. Shelkov,[3] T. Skwarnicki,[3] R. Stroynowski,[3] I. Volobouev,[3] G. Wei,[3] M. Artuso,[4] M. Gao,[4] M. Goldberg,[4] D. He,[4] N. Horwitz,[4] S. Kopp,[4] G.C. Moneti,[4] R. Mountain,[4] F. Muheim,[4] Y. Mukhin,[4] S. Playfer,[4] S. Stone,[4] X. Xing,[4] J. Bartelt,[5] S.E. Csorna,[5] V. Jain,[5] S. Marka,[5] D. Gibaut,[6] K. Kinoshita,[6] P. Pomianowski,[6] B. Barish,[7] M. Chadha,[7] S. Chan,[7] D.F. Cowen,[7] G. Eigen,[7] J.S. Miller,[7] C. O'Grady,[7] J. Urheim,[7] A.J. Weinstein,[7] F. Würthwein,[7] D.M. Asner,[8] M. Athanas,[8] D.W. Bliss,[8] W.S. Brower,[8] G. Masek,[8] H.P. Paar,[8] J. Gronberg,[9] C.M. Korte,[9] R. Kutschke,[9] S. Menary,[9] R.J. Morrison,[9] S. Nakanishi,[9] H.N. Nelson,[9] T.K. Nelson,[9] C. Qiao,[9] J.D. Richman,[9] D. Roberts,[9] A. Ryd,[9] H. Tajima,[9] M.S. Witherell,[9] R. Balest,[10] K. Cho,[10] W.T. Ford,[10] M. Lohner,[10] H. Park,[10] P. Rankin,[10] J.G. Smith,[10] J.P. Alexander,[11] C. Bebek,[11] B.E. Berger,[11] K. Berkelman,[11] K. Bloom,[11] T.E. Browder,[11*] D.G. Cassel,[11] H.A. Cho,[11] D.M. Coffman,[11] D.S. Crowcroft,[11] M. Dickson,[11] P.S. Drell,[11] D.J. Dumas,[11] R. Ehrlich,[11] R. Elia,[11] P. Gaidarev,[11] M. Garcia-Sciveres,[11] B. Gittelman,[11] S.W. Gray,[11] D.L. Hartill,[11] B.K. Heltsley,[11] S. Henderson,[11] C.D. Jones,[11] S.L. Jones,[11] J. Kandaswamy,[11] N. Katayama,[11] P.C. Kim,[11] D.L. Kreinick,[11] T. Lee,[11] Y. Liu,[11] G.S. Ludwig,[11] J. Masui,[11] J. Mevissen,[11] N.B. Mistry,[11] C.R. Ng,[11] E. Nordberg,[11] J.R. Patterson,[11] D. Peterson,[11] D. Riley,[11] A. Soffer,[11] P. Avery,[12] A. Freyberger,[12] K. Lingel,[12] C. Prescott,[12] J. Rodriguez,[12] S. Yang,[12] J. Yelton,[12] G. Brandenburg,[13] D. Cinabro,[13] T. Liu,[13] M. Saulnier,[13] R. Wilson,[13] H. Yamamoto,[13] T. Bergfeld,[14] B.I. Eisenstein,[14] J. Ernst,[14] G.E. Gladding,[14] G.D. Gollin,[14] M. Palmer,[14] M. Selen,[14] J.J. Thaler,[14] K.W. Edwards,[15] K.W. McLean,[15] M. Ogg,[15] A. Bellerive,[16] D.I. Britton,[16] E.R.F. Hyatt,[16] R. Janicek,[16] D.B. MacFarlane,[16] P.M. Patel,[16] B. Spaan,[16] A.J. Sadoff,[17] R. Ammar,[18] P. Baringer,[18] A. Bean,[18] D. Besson,[18] D. Coppage,[18] N. Copty,[18] R. Davis,[18] N. Hancock,[18] S. Kotov,[18] I. Kravchenko,[18] N. Kwak,[18] Y. Kubota,[19] M. Lattery,[19]

M. Momayezi,[19] J.K. Nelson,[19] S. Patton,[19] R. Poling,[19] V. Savinov,[19] S. Schrenk,[19]
R. Wang,[19] M.S. Alam,[20] I.J. Kim,[20] Z. Ling,[20] A.H. Mahmood,[20] J.J. O'Neill,[20]
H. Severini,[20] C.R. Sun,[20] F. Wappler,[20] G. Crawford,[21] R. Fulton,[21] D. Fujino,[21]
K.K. Gan,[21] K. Honscheid,[21] H. Kagan,[21] R. Kass,[21] J. Lee,[21] M. Sung,[21] C. White,[21]
A. Wolf,[21] M.M. Zoeller,[21] X. Fu,[22] B. Nemati,[22] W.R. Ross,[22] P. Skubic,[22] and M. Wood[22]

(CLEO Collaboration)

[1] *Purdue University, West Lafayette, Indiana 47907*
[2] *University of Rochester, Rochester, New York 14627*
[3] *Southern Methodist University, Dallas, Texas 75275*
[4] *Syracuse University, Syracuse, New York 13244*
[5] *Vanderbilt University, Nashville, Tennessee 37235*
[6] *Virginia Polytechnic Institute and State University, Blacksburg, Virginia, 24061*
[7] *California Institute of Technology, Pasadena, California 91125*
[8] *University of California, San Diego, La Jolla, California 92093*
[9] *University of California, Santa Barbara, California 93106*
[10] *University of Colorado, Boulder, Colorado 80309-0390*
[11] *Cornell University, Ithaca, New York 14853*
[12] *University of Florida, Gainesville, Florida 32611*
[13] *Harvard University, Cambridge, Massachusetts 02138*
[14] *University of Illinois, Champaign-Urbana, Illinois, 61801*
[15] *Carleton University, Ottawa, Ontario K1S 5B6 and the Institute of Particle Physics, Canada*
[16] *McGill University, Montréal, Québec H3A 2T8 and the Institute of Particle Physics, Canada*
[17] *Ithaca College, Ithaca, New York 14850*
[18] *University of Kansas, Lawrence, Kansas 66045*
[19] *University of Minnesota, Minneapolis, Minnesota 55455*
[20] *State University of New York at Albany, Albany, New York 12222*
[21] *Ohio State University, Columbus, Ohio, 43210*
[22] *University of Oklahoma, Norman, Oklahoma 73019*

(July 7, 1995)

Abstract

We have observed two new decay modes of the charmed baryon Ξ_c^+ into $\Sigma^+K^-\pi^+$ and Σ^+K^{*0} using data collected with the CLEO II detector. We also present the first measurement of the branching fraction for the previously observed decay mode $\Xi_c^+ \to \Lambda K^-\pi^+\pi^+$. The branching fractions for these three modes relative to $\Xi_c^+ \to \Xi^-\pi^+\pi^+$ are measured to be $1.18 \pm 0.26 \pm 0.17$, $0.92 \pm 0.27 \pm 0.14$, and $0.58 \pm 0.16 \pm 0.07$, respectively.

---

*Permanent address: University of Hawaii at Manoa

I. INTRODUCTION

The Ξ_c^+ charmed baryon lifetime is $2.5 - 3.5$ times longer than the Ξ_c^0 lifetime [1,2]. It is widely believed that destructive interference between the external and internal W-emission diagrams in Ξ_c^+ decays and the presence of W-exchange decay channels in Ξ_c^0 decays contribute to the difference in lifetimes [3]. Measuring the branching fractions of Ξ_c decay modes can help us determine the mechanisms for Ξ_c charmed baryon decays and verify these assertions.

Although numerous decay modes of the Λ_c^+ charmed baryon have been observed, only a few Ξ_c^+ decay modes have been detected. The $\Xi_c^+ \to \Xi^-\pi^+\pi^+$ and $\Lambda K^-\pi^+\pi^+$ decay modes [4] have been previously observed by a number of groups [1,5]. CLEO has also observed the Ξ_c^+ decaying into modes with a neutral Ξ^0 hyperon, namely, $\Xi_c^+ \to \Xi^0\pi^+$, $\Xi^0\pi^+\pi^0$, and $\Xi^0\pi^+\pi^-\pi^+$ [6]. The simplest external W spectator diagram produces a Ξ^0 in the final state. Other hyperons (Λ, Σ^+, and Ξ^-) can be produced either through internal W-emission diagrams or by combining valence quarks with quarks popped from the vacuum.

We report here the observation of two new decay modes of the Ξ_c^+ into $\Sigma^+K^-\pi^+$ and Σ^+K^{*0}. The simplest diagram yielding Σ^+K^{*0} is an internal W-emission diagram. The $\Sigma^+K^-\pi^+$ mode has been previously seen by the CERN fixed target experiment ACCMOR [7], but is based on three events in the mass range $2445 - 2475$ MeV/c^2. We also present a first measurement of the $\Xi_c^+ \to \Lambda K^-\pi^+\pi^+$ branching fraction.

The data were collected with the CLEO II detector at the Cornell e^+e^- storage ring CESR, which operated on, and just below, the $\Upsilon(4S)$ resonance. The CLEO II detector [8] is a large solenoidal detector with 67 tracking layers and a CsI electromagnetic calorimeter that provides efficient π^0 reconstruction. We have used a total integrated luminosity of 3.6 fb^{-1}, which corresponds to roughly 4 million $c\bar{c}$ events.

II. PARTICLE ID

Charged proton and pion candidates are required to have specific ionization loss (dE/dx) information and, when available, time-of-flight information consistent with the value expected for the assumed particle type. We use clean samples of $\Lambda \to p\pi^-$, $D^0 \to K^-\pi^+$ from D^{*+}'s, and $K_s^0 \to \pi^+\pi^-$ to measure the particle identification efficiencies directly from the data for protons, kaons, and pions, respectively.

The proton in our Ξ_c^+ decay modes always comes from a hyperon (Λ, Σ^+, or Ξ^-) and backgrounds from other hadrons are relatively low. Charged kaons, however, are selected

with stronger particle identification cuts to minimize the reflection peaks in the Ξ_c^+ invariant mass distributions from the $\Lambda_c^+ \to \Lambda\pi^+\pi^-\pi^+$ and $\Sigma^+\pi^-\pi^+$ decay modes, where one of the pions is mis-identified as a kaon. These reflections will produce broad distributions peaked near the Ξ_c^+ mass. We require the probability that the candidate is a kaon to be at least 50% of the sum of the probabilities for the proton, kaon, and pion hypotheses.

III. SELECTION OF HYPERONS

The Λ candidates are selected through their decay $\Lambda \to p\pi^-$ by reconstructing a secondary decay vertex from the intersection of two oppositely charged tracks in the plane perpendicular to the beam axis (the $r-\phi$ plane). Similarly, Ξ^- hyperons are selected through the decay mode $\Xi^- \to \Lambda\pi^-$ by reconstructing a secondary decay vertex from the intersection of a Λ candidate and a π^- particle in the $r-\phi$ plane. The Σ^+ hyperon candidates are selected from $p\pi^0$ combinations that are consistent with coming from a decay vertex displaced from the primary interaction point [9]. The invariant masses of the Λ, Ξ^-, and Σ^+ candidates must lie within 5, 6, and 15 MeV/c^2 of their nominal values ($\sim 3\sigma$), respectively.

IV. OBSERVATION OF $\Xi_c^+ \to \Sigma^+K^-\pi^+$ AND Σ^+K^{*0}

Charmed baryons from $e^+e^- \to c\bar{c}$ interactions are produced with a hard momentum spectrum, so we reduce the combinatoric background by requiring either $x_p > 0.5$ or $x_p > 0.6$, depending on the decay mode, where $x_p = P_{\Xi_c^+}/\sqrt{E_{beam}^2 - M_{\Xi_c^+}^2}$ is the scaled momentum of the Ξ_c^+. The x_p cut also eliminates Ξ_c^+ baryons that arise from B meson decays. In addition, we require that the daughter particles of the Ξ_c^+ lie within 90 degrees of the candidate Ξ_c^+ momentum vector.

The invariant mass distribution for $\Xi_c^+ \to \Sigma^+K^-\pi^+$ candidates with $x_p > 0.5$ is shown in Figure 1. We observe a Ξ_c^+ signal at $\sim$ 2470 MeV/c^2. However, the distribution also contains a reflection peak from $\Lambda_c^+ \to \Sigma^+\pi^-\pi^+$ events where a pion is misidentified as a kaon. The shape of the reflection peak is determined from a Monte Carlo phase space decay of $\Lambda_c^+ \to \Sigma^+\pi^-\pi^+$. The area is determined from data by constructing the invariant mass distribution for the $\Sigma^+\pi^-\pi^+$ hypothesis from all $\Sigma^+K^-\pi^+$ combinations in Figure 1. A $\Lambda_c^+ \to \Sigma^+\pi^-\pi^+$ signal of 142 ± 25 events is observed, which is taken to be the area of the reflection peak.

We parametrize the $\Sigma^+K^-\pi^+$ mass distribution by a Gaussian signal, the $\Lambda_c^+ \to \Sigma^+\pi^-\pi^+$ reflection peak with fixed area of 142 events, and a 3rd order Chebyshev polynomial back-



FIG. 1. Invariant mass distribution for $\Xi_c^+ \to \Sigma^+ K^- \pi^+$. The curve at the bottom corresponds to the reflection peak from misidentified $\Lambda_c^+ \to \Sigma^+ \pi^- \pi^+$ events.

ground. The width of the Gaussian is taken from Monte Carlo studies to be $\sigma = 8.6$ MeV/c^2. We observe 119 ± 23 events (statistical error only) in the $\Xi_c^+ \to \Sigma^+ K^- \pi^+$ decay mode at a mass of 2469.9 ± 2.0 MeV/c^2 (statistical error only), consistent with the nominal Ξ_c^+ mass. Excluding the Λ_c^+ reflection in the fit artificially raises the Ξ_c^+ yield.

A search was made for the two-body decay $\Xi_c^+ \to \Sigma^+ K^{*0}$ by examining the resonant substructure of the $\Sigma^+ K^- \pi^+$ mode. We divide the $K^- \pi^+$ mass range into seven regions (of unequal size) from 0.6 to 1.3 GeV/c^2 and obtain the Ξ_c^+ yield for each region. Contributions from the Λ_c^+ reflection are again measured and incorporated in the fit. Figure 2 shows a clear K^{*0} signal in the $K^- \pi^+$ invariant mass distribution. We fit the data (points) to the sum of two curves representing the resonant and non-resonant (n.r.) $\Sigma^+ K^- \pi^+$ contributions (histogram), where the shapes but not the normalizations are determined from Monte Carlo. The fit yields 61 ± 17 events from $\Xi_c^+ \to \Sigma^+ K^{*0}$ and 55 ± 22 events from n.r. $\Xi_c^+ \to \Sigma^+ K^- \pi^+$ decays. The efficiency corrected yields are 619 ± 172 events for the $\Xi_c^+ \to \Sigma^+ K^{*0}$, $K^{*0} \to K^- \pi^+$ channel and 525 ± 208 events for the n.r. $\Xi_c^+ \to \Sigma^+ K^- \pi^+$ mode.

As a check, we also measure the $\Xi_c^+ \to \Sigma^+ K^{*0}$ contribution by examining the $\Sigma^+ K^- \pi^+$ invariant mass distribution after requiring the $K^- \pi^+$ mass to be within 50 MeV/c^2 of the K^{*0} mass (see Figure 3). About half the Ξ_c^+ signal remains and the backgrounds are greatly reduced. A fit to the $\Sigma^+ K^- \pi^+$ mass distribution yields 59 ± 12 events. We have included the $\Sigma^+ \pi^- \pi^+$ reflection peak with a fixed area of 35 events, as determined in the data. Since the K^{*0} mass cut is wide, a non-negligible amount of non-resonant $\Xi_c^+ \to \Sigma^+ K^- \pi^+$ decays



FIG. 2. The $K^-\pi^+$ invariant mass distribution for the $\Xi_c^+ \to \Sigma^+ K^-\pi^+$ decay mode. The points are from data, where the fitted $\Xi_c^+ \to \Sigma^+ K^-\pi^+$ yield is plotted for each bin of $K^-\pi^+$ mass. The histogram is a χ^2 fit to the sum of two curves representing the Monte Carlo shapes for resonant $\Sigma^+ K^{*0}$ and non-resonant $\Sigma^+ K^-\pi^+$.



FIG. 3. Invariant mass distribution for $\Xi_c^+ \to \Sigma^+ K^-\pi^+$ with the K^{*0} mass cut of $|M_{K\pi} - 892| < 50$ MeV/c^2. The curve at the bottom corresponds to the reflection peak from misidentified $\Lambda_c^+ \to \Sigma^+\pi^-\pi^+$ events.



FIG. 4. Invariant mass distribution for $\Xi_c^+ \to \Lambda K^-\pi^+\pi^+$. The curve at the bottom corresponds to the reflection peak from misidentified $\Lambda_c^+ \to \Lambda\pi^+\pi^-\pi^+$ events.

can survive the cut, although the reconstruction efficiency is a factor of four smaller than that of resonant Σ^+K^{*0}. We obtain a nearly pure sample of n.r. $\Sigma^+K^-\pi^+$ by requiring that $|M_{K\pi} - 892| > 100$ MeV/c^2. In this case, we measure 45 ± 17 events in the data. We unfold the resonant and n.r. contributions from these two measurements and obtain an efficiency corrected yield of 669 ± 179 events for the decay chain $\Xi_c^+ \to \Sigma^+K^{*0}$, $K^{*0} \to K^-\pi^+$ and 550 ± 252 events for the non-resonant mode $\Xi_c^+ \to \Sigma^+K^-\pi^+$. This is consistent with the first method and the difference is taken as a systematic error.

V. OBSERVATION OF $\Xi_c^+ \to \Lambda K^-\pi^+\pi^+$

The invariant mass distribution for $\Xi_c^+ \to \Lambda K^-\pi^+\pi^+$ candidates with $x_p > 0.6$ is shown in Figure 4. The more stringent x_p requirement is needed to reduce the combinatoric background for this higher multiplicity final state. We parametrize the mass distribution as the sum of a Gaussian signal, a contribution from the $\Lambda_c^+ \to \Lambda\pi^+\pi^-\pi^+$ reflection peak, and a 3rd order Chebyshev polynomial background. The width of the Gaussian is taken from Monte Carlo studies to be $\sigma = 7.0$ MeV/c^2. The reflection peak is fixed to the $\Lambda_c^+ \to \Lambda\pi^+\pi^-\pi^+$ yield of 116 ± 22 events, determined from a fit to the invariant mass distribution for the $\Lambda\pi^+\pi^-\pi^+$ hypothesis for all $\Lambda K^-\pi^+\pi^+$ combinations in Figure 4.

We observe 61 ± 15 events for the mode $\Xi_c^+ \to \Lambda K^-\pi^+\pi^+$ at a mass of 2467.5 ± 2.0 MeV/c^2. We have also looked for possible K^{*0} and $\Sigma^{*+}(1385)$ resonant substructure in $\Xi_c^+ \to \Lambda K^-\pi^+\pi^+$ decays. None is observed, and we place upper limits of



FIG. 5. Invariant mass distribution for $\Xi_c^+ \to \Xi^-\pi^+\pi^+$.

$\mathcal{B}(\Lambda K^{*0}\pi^+)/\mathcal{B}(\Lambda K^-\pi^+\pi^+) < 0.5$ and $\mathcal{B}(\Sigma^{*+}K^-\pi^+)/\mathcal{B}(\Lambda K^-\pi^+\pi^+) < 0.7$ at the 90% confidence level.

VI. THE $\Xi_c^+ \to \Xi^-\pi^+\pi^+$ DECAY MODE

Since the Ξ_c^+ total cross section is not precisely known, we convert our observations into branching fractions relative to the well measured decay mode $\Xi_c^+ \to \Xi^-\pi^+\pi^+$. For each mode we apply the same Ξ_c^+ momentum cut to $\Xi^-\pi^+\pi^+$ (either 0.5 or 0.6) to reduce the systematic dependence on the x_p cut.

The invariant mass distribution for $\Xi_c^+ \to \Xi^-\pi^+\pi^+$ candidates with $x_p > 0.5$ is shown in Figure 5. We parametrize the mass distribution by a Gaussian signal and a quadratic polynomial background. The width of the Gaussian is taken from Monte Carlo studies to be $\sigma = 10.1$ MeV/c^2. We observe 131 ± 14 events at a mass of 2466.6 ± 1.3 MeV/c^2. For $x_p > 0.6$ we observe 100 ± 11 Ξ_c^+ events.

No resonant substructure is observed, and we place an upper limit for the two-body decay $\Xi_c^+ \to \Xi_c^{*0}(1530)\pi^+$ to be $\mathcal{B}(\Xi_c^+ \to \Xi_c^{*0}\pi^+)/\mathcal{B}(\Xi_c^+ \to \Xi^-\pi^+\pi^+) < 0.2$ at the 90% confidence level. Körner and Krämer [10] have predicted that this decay rate should be zero.

VII. DISCUSSION

The raw yields, efficiencies, and resultant branching fractions for all decay modes are shown in Table I. The systematic errors associated with the measured relative branching

TABLE I. Summary of results on new Ξ_c^+ decay modes. The $\Sigma^+K^-\pi^+$ mode includes both resonant and non-resonant contributions. The efficiencies ($\mathcal{E}$) do not include branching fractions to the observed final states. The first error in the relative branching fraction is statistical and the second is systematic.

Decay Mode	x_p cut	Events	$\mathcal{E}$ (%)	$\mathcal{B}/\mathcal{B}(\Xi_c^+ \to \Xi^-\pi^+\pi^+)$
$\Sigma^+K^-\pi^+$	0.5	119 ± 23	10.4	$1.18 \pm 0.26 \pm 0.17$
Σ^+K^{*0}	0.5	61 ± 17	9.8	$0.92 \pm 0.27 \pm 0.14$
$\Lambda K^-\pi^+\pi^+$	0.6	61 ± 15	11.5	$0.58 \pm 0.16 \pm 0.07$
$\Xi^-\pi^+\pi^+$	0.5	131 ± 14	10.6	1.0

TABLE II. Summary of systematic errors.

Source	Error (%)		
	$\Sigma^+K^-\pi^+$	Σ^+K^{*0}	$\Lambda K^-\pi^+\pi^+$
Λ and Σ^+ finding	6	6	5
Ξ^- finding	7	7	7
π^0 finding	5	5	—
tracking	4	4	—
kaon and proton identification	5	5	5
float vs fixed $\sigma(\Xi_c^+)$	6	3	2
area of Λ_c^+ reflection peak	3	2	5
shape of Λ_c^+ reflection peak	—	—	3
K^{*0} fit method	—	8	—
TOTAL	14	15	12]

fractions of the Ξ_c^+ decay modes are listed in Table II. The main sources of systematic error are due to uncertainties in the efficiencies for Λ, Σ^+, and Ξ^- reconstruction ($5-7\%$), particle identification (5%), charged particle tracking (4%), and π^0 reconstruction (5%). An additional 8% error is assigned to the Σ^+K^{*0} yield for the difference between the two methods of extracting the K^{*0} contribution. Variations in the Ξ_c^+ yields due to uncertainties in the signal width are $2-6\%$ and in the area of the reflection peak for $\Lambda_c^+ \to \Lambda\pi^+\pi^-\pi^+$ and $\Sigma^+\pi^-\pi^+$ are $2-5\%$, depending on the channel. CLEO observes that about half of the $\Lambda_c^+ \to \Lambda\pi^+\pi^-\pi^+$ decays proceed via a Σ^{*+} or Σ^{*-} resonance [11]. Variations in the shape of the Λ_c^+ reflection peak due to the $\Sigma^{*\pm}$ substructure correspond to a 3% systematic error. The total systematic error for the $\Sigma^+K^-\pi^+$, Σ^+K^{*0}, and $\Lambda K^-\pi^+\pi^+$ decay modes are 14%, 15%, and 12%, respectively.

Körner and Krämer [10], Zenczykowski [12], and Datta [13] have used quark and pole models to make theoretical predictions for the Ξ_c^+ decay into the two-body final state $\Sigma^+ K^{*0}$. They calculate $\Gamma(\Sigma^+ K^{*0})$ to be $5.3 \times 10^{10} s^{-1}$, $8.7 \times 10^{10} s^{-1}$, and $3.6 \times 10^{10} s^{-1}$, respectively. We can convert our measured branching fraction into a decay rate by using the world average Ξ_c^+ lifetime [1] and CLEO's measurement of $\mathcal{B}(\Xi_c^+ \to \Xi^- \pi^+ \pi^+) = f_{\Xi_c}(2.1 \pm 0.8 \pm 0.4)\%$ [2], where the fraction $f_{\Xi_c} \equiv \mathcal{B}(\Xi_c^+ \to \Xi^0 \ell \nu)/\mathcal{B}(\Xi_c^+ \to \ell X)$ is expected to be in the range $0.6 - 1.0$. We thus obtain $\Gamma(\Sigma^+ K^{*0}) = f_{\Xi_c}(5.5 \pm 2.9) \times 10^{10} s^{-1}$, in good agreement with the theoretical predictions.

In summary, we have observed two new decay modes of the Ξ_c^+ into $\Sigma^+ K^- \pi^+$ and $\Sigma^+ K^{*0}$. We also have a first measurement of the relative branching fraction for $\Xi_c^+ \to \Lambda K^- \pi^+ \pi^+$. The ratios of branching fractions suggest that the two resultant strange quarks do not always combine to form a Ξ hyperon, but can end up in two separate hadrons via an internal W-emission diagram. Our measurement of $\mathcal{B}(\Lambda K^- \pi^+ \pi^+)/\mathcal{B}(\Xi^- \pi^+ \pi^+)$ is considerably lower than a preliminary estimate by the WA89 experiment of ≈ 4 [5]. Somewhat surprisingly, neither $\Lambda K^- \pi^+ \pi^+$ or $\Xi^- \pi^+ \pi^+$ have any observable resonant substructure. This behavior is similar to the Λ_c^+ charmed baryon in which only a quarter of the observed Λ_c^+ hadronic width (about 40% of the width has been accounted for) goes through two-body decay modes, with the bulk proceeding most likely through multibody states. In contrast, about half of the hadronic width of D^+ and D^0 mesons proceeds via two-body decays. Clearly, charmed baryon decays continue to be an interesting laboratory for weak decay physics.

ACKNOWLEDGEMENTS

We gratefully acknowledge the effort of the CESR staff in providing us with excellent luminosity and running conditions. This work was supported by the National Science Foundation, the U.S. Department of Energy, the Heisenberg Foundation, the Alexander von Humboldt Stiftung, the Natural Sciences and Engineering Research Council of Canada, and the A.P. Sloan Foundation.

REFERENCES

[1] Particle Data Group, L. Montanet *et al.*, Phys. Rev. D **50**, 1173 (1994).

[2] J.P. Alexander *et al.* (CLEO Collab.), Phys. Rev. Lett. **74**, 3113 (1995), and an erratum to appear in PRL (1995).

[3] B. Guberina, R. Rückl, and J. Trampetić, Z. Phys. C **33**, 297 (1986); M.B. Voloshin and M.A. Shifman, Sov. Phys. JETP **64**, 698 (1986).

[4] Unless otherwise stated, the charge conjugate mode is always implied.

[5] E. Chudakov (WA89 Collab.), "Charmed Baryon Production in the CERN Hyperon Beam," Heavy Quark 94 Conf., Virginia, Oct. 1994.

[6] A. Bean *et al.* (CLEO Collab.), CLEO CONF 93-8.

[7] S. Barlag *et al.* (ACCMOR Collab.), Phys. Lett. B **233**, 522 (1989).

[8] Y. Kubota *et al.* (CLEO Collab.), Nucl. Inst. and Meth. **A320**, 66 (1992).

[9] Y. Kubota *et al.* (CLEO Collab.), Phys. Rev. Lett. **71**, 3255 (1993).

[10] J.G. Körner and M. Krämer, Z. Phys. C **55**, 659 (1992).

[11] CLEO internal note, CBX-92-61.

[12] P. Zenczykowski, Phys. Rev. D **50**, 402 (1994). The value for $\mathcal{B}(\Sigma^+ K^{*0})$ used the Ξ_c^+ lifetime from the 1990 PDG, which we have corrected to reflect the newer 1994 PDG average.

[13] A. Datta, "Non-Leptonic two body decays of Charmed and Λ_b Baryons," preprint UH-511-824-95.

[14] A.V. Monohar and M.B. Wise, Phys. Rev. D **49**, 1310 (1994).