

Dalitz Analysis of $D^+ \rightarrow \pi^+ \pi^- \pi^{+*}$

G. Bonvicini,¹ D. Cinabro,¹ M. Dubrovin,¹ A. Lincoln,¹ D. M. Asner,² K. W. Edwards,² R. A. Briere,³ I. Brock,³ J. Chen,³ T. Ferguson,³ G. Tatishvili,³ H. Vogel,³ M. E. Watkins,³ J. L. Rosner,⁴ N. E. Adam,⁵ J. P. Alexander,⁵ K. Berkelman,⁵ D. G. Cassel,⁵ J. E. Duboscq,⁵ K. M. Ecklund,⁵ R. Ehrlich,⁵ L. Fields,⁵ L. Gibbons,⁵ R. Gray,⁵ S. W. Gray,⁵ D. L. Hartill,⁵ B. K. Heltsley,⁵ D. Hertz,⁵ C. D. Jones,⁵ J. Kandaswamy,⁵ D. L. Kreinick,⁵ V. E. Kuznetsov,⁵ H. Mahlke-Krüger,⁵ P. U. E. Onyisi,⁵ J. R. Patterson,⁵ D. Peterson,⁵ J. Pivarski,⁵ D. Riley,⁵ A. Ryd,⁵ A. J. Sadoff,⁵ H. Schwarhoff,⁵ X. Shi,⁵ S. Stroiney,⁵ W. M. Sun,⁵ T. Wilksen,⁵ M. Weinberger,⁵ S. B. Athar,⁶ R. Patel,⁶ V. Potlia,⁶ J. Yelton,⁶ P. Rubin,⁷ C. Cawfield,⁸ B. I. Eisenstein,⁸ I. Karliner,⁸ D. Kim,⁸ N. Lowrey,⁸ P. Naik,⁸ C. Sedlack,⁸ M. Selen,⁸ E. J. White,⁸ J. Wiss,⁸ M. R. Shepherd,⁹ D. Besson,¹⁰ T. K. Pedlar,¹¹ D. Cronin-Hennessy,¹² K. Y. Gao,¹² D. T. Gong,¹² J. Hietala,¹² Y. Kubota,¹² T. Klein,¹² B. W. Lang,¹² R. Poling,¹² A. W. Scott,¹² A. Smith,¹² P. Zweber,¹² S. Dobbs,¹³ Z. Metreveli,¹³ K. K. Seth,¹³ A. Tomaradze,¹³ J. Ernst,¹⁴ H. Severini,¹⁵ S. A. Dytman,¹⁶ W. Love,¹⁶ V. Savinov,¹⁶ O. Aquines,¹⁷ Z. Li,¹⁷ A. Lopez,¹⁷ S. Mehrabyan,¹⁷ H. Mendez,¹⁷ J. Ramirez,¹⁷ G. S. Huang,¹⁸ D. H. Miller,¹⁸ V. Pavlunin,¹⁸ B. Sanghi,¹⁸ I. P. J. Shipsey,¹⁸ B. Xin,¹⁸ G. S. Adams,¹⁹ M. Anderson,¹⁹ J. P. Cummings,¹⁹ I. Danko,¹⁹ J. Napolitano,¹⁹ Q. He,²⁰ J. Insler,²⁰ H. Muramatsu,²⁰ C. S. Park,²⁰ E. H. Thorndike,²⁰ F. Yang,²⁰ T. E. Coan,²¹ Y. S. Gao,²¹ F. Liu,²¹ M. Artuso,²² S. Blusk,²² J. Butt,²² J. Li,²² N. Menaa,²² R. Mountain,²² S. Nisar,²² K. Randrianarivony,²² R. Redjimi,²² R. Sia,²² T. Skwarnicki,²² S. Stone,²² J. C. Wang,²² K. Zhang,²² and S. E. Csorna²³

(CLEO Collaboration)

¹Wayne State University, Detroit, Michigan 48202

²Carleton University, Ottawa, Ontario, Canada K1S 5B6

³Carnegie Mellon University, Pittsburgh, Pennsylvania 15213

⁴Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637

⁵Cornell University, Ithaca, New York 14853

⁶University of Florida, Gainesville, Florida 32611

⁷George Mason University, Fairfax, Virginia 22030

⁸University of Illinois, Urbana-Champaign, Illinois 61801

⁹Indiana University, Bloomington, Indiana 47405

¹⁰University of Kansas, Lawrence, Kansas 66045

¹¹Luther College, Decorah, Iowa 52101

¹²University of Minnesota, Minneapolis, Minnesota 55455

¹³Northwestern University, Evanston, Illinois 60208

¹⁴State University of New York at Albany, Albany, New York 12222

¹⁵University of Oklahoma, Norman, Oklahoma 73019

¹⁶University of Pittsburgh, Pittsburgh, Pennsylvania 15260

¹⁷University of Puerto Rico, Mayaguez, Puerto Rico 00681

¹⁸Purdue University, West Lafayette, Indiana 47907

¹⁹Rensselaer Polytechnic Institute, Troy, New York 12180

²⁰University of Rochester, Rochester, New York 14627

²¹Southern Methodist University, Dallas, Texas 75275

²²*Syracuse University, Syracuse, New York 13244*
²³*Vanderbilt University, Nashville, Tennessee 37235*
(Dated: July 24, 2006)

Abstract

Using 281/pb of data recorded by the CLEO-c detector observing e^+e^- collisions at the $\psi(3770)$, corresponding to 1.8 million $D\bar{D}$ pairs, the substructure of the decay $D^+ \rightarrow \pi^+\pi^-\pi^+$ is investigated using the Dalitz plot technique. The results presented in this document are preliminary.

*Submitted to the 33rd International Conference on High Energy Physics, July 26 - August 2, 2006, Moscow

A Dalitz plot analysis [1] of $D^+ \rightarrow \pi^+\pi^-\pi^+$ has previously been done by E791 [2] and FOCUS [3]. The preliminary analysis described here is from CLEO-c [4], and represents the first time we have done the same Dalitz plot analysis as the fixed target experiments. Previously CLEO has focused on analyses with π^0 's in the final state. The decay is selected with cuts on the beam constrained mass of three charged tracks consistent with pions and the difference of their energy from the beam energy. A sample of 4100 events is selected with a signal to noise of about two to one. The E791 and FOCUS samples are of similar size and cleanliness.

The Dalitz plot is symmetric under the interchange of like-sign pions thus we do the analysis in the two dimensions of high unlike-sign pion mass squared versus low unlike-sign pion mass. There is a large contribution from $K_S\pi$ which because of the long K_S lifetime should not interfere with the other contributions to the plot. We do not consider this stripe on the Dalitz plot when fitting for two body resonance contributions. The efficiency across the Dalitz plot is modeled with simulated events that are fit to a two-dimensional second order polynomial. While there is a notable fall of the efficiency in the corners of the Dalitz plot the changes are smooth, and well modeled by the polynomial. Backgrounds are taken from sidebands and extra resonance contributions to the background are allowed from mismeasured K_S , ρ , and $f_2(1270)$ decays. Many possible resonances can contribute to the decay, and a total of 13 different resonances are considered. Parameters describing these resonances are taken from previous experiments. Only contributions with an amplitude significant at more than three standard deviations are said to be observed, and others are limited. Contributions that are not significant are not included in the decay model used for the result.

Figure 1 shows the Dalitz plot and projections on to the squared masses. Contributions from $\rho^0\pi$ and $f_2(1270)\pi$ are clearly visible. Table I shows the preliminary fit fractions measured by CLEO comparing with the results of the E791 analysis mentioned above. There

TABLE I: Comparison of the fit fractions, in percent, found in the Dalitz plot analysis of $D^+ \rightarrow \pi^+\pi^-\pi^+$ between the preliminary CLEO analysis and E791. CLEO limits are at 90% confidence level.

Contribution	CLEO	E791
$\rho^0\pi^+$	20.0 ± 2.5	33.6 ± 3.9
$\sigma^0\pi^+$	41.8 ± 2.9	46.3 ± 9.2
$f_2(1270)\pi^+$	18.2 ± 2.7	19.4 ± 2.5
$f_0(980)\pi^+$	4.1 ± 0.9	6.2 ± 1.4
$f_0(1370)\pi^+$	2.6 ± 1.9	2.3 ± 1.7
$f_0(1500)\pi^+$	3.4 ± 1.3	
Non-resonant	< 3.5	7.8 ± 6.6
$\rho(1450)\pi^+$	< 2.4	0.7 ± 0.8

is broad agreement between the two results, including the observation of a $\sigma\pi$ contribution. In an alternative analysis using the same decay model as E791 the agreement is only slightly better, but the fit is much less likely to model our data than the model shown in the table which does not include non-resonant and $\rho(1450)\pi^+$ contributions, but does include a $f_0(1500)\pi^+$ contribution. Models without a $\sigma\pi$ contribution do not agree well with the data.




FIG. 1: Dalitz plot and projections for $D^+ \rightarrow \pi^+ \pi^- \pi^+$

This CLEO analysis is preliminary, and we plan to consider a generalized model of $\pi\pi$ S-wave interactions to model σ and f_0 contributions such as the K-matrix which is used in the FOCUS analysis mentioned above.

We gratefully acknowledge the effort of the CESR staff in providing us with excellent luminosity and running conditions. D. Cronin-Hennessy and A. Ryd thank the A.P. Sloan Foundation. This work was supported by the National Science Foundation, the U.S. Department of Energy, and the Natural Sciences and Engineering Research Council of Canada.

-
- [1] R.H.Dalitz, Phil. Mag. 44 (1953) 1068.
 [2] E.M.Aitala *et al.* (Fermilab E791 Collaboration), Phys. Rev. Lett. 86 (2001) 770.
 [3] J.M.Link *et al.* (FOCUS Collaboration), Phys. Lett. B 585 (2004) 200.
 [4] R.A. Briere *et al.* (CESR-c and CLEO-c Taskforces, CLEO-c Collaboration), Cornell University, LEPP Report No. CLNS 01/1742 (2001) (unpublished).