


UFRJ



Universidade Federal  
do Rio de Janeiro



Sensacional!

# The Physics of Charm: Recent Experimental Results


Jim Napolitano  
Rensselaer Polytechnic Institute  
*and the CLEO-c Collaboration*

XXVI Physics in Collision 6-9 July 2006 Buzios, Brazil




Rensselaer

CLEO-c



# What is *Special* about Charm?


1) *Its mass*. The charm quark is “heavy”...



... but not “too” heavy.

*Most decay modes of hadrons with charm are “easy” to observe experimentally.*

2) *Its decays.* Charm is the only heavy quark that forms hadrons with CKM-allowed decays.




# The Experiments


$e^+e^-$  and Photon Beams



# BABAR





# BES



# CLEO

Hadroproduction




# E835

# SELEX



# A Survey of Recent Results

*My own selection! My apologies for not covering it all!*

- “Precision” Measurements  
Small error bars, stringent limits, and tying up old loose ends
- New States *(Note: Talk tomorrow by R. Waldi)*  
A resurgence in charmonium
- Confronting Lattice QCD  
Testing “high precision” lattice calculations
- $D^0$  Mixing and Tests of CP Violation  
Current status and future prospects



# The $D_s$ Lifetime




$$\gamma N \rightarrow (K^+ K^- \pi^\pm) X$$



$D_s$  signal

Sidebands

$$\tau(D_s) = 507.4 \pm 5.5 \pm 5.1 \text{ fs}$$




# Limit: $D^+ \rightarrow \pi^+ \mu^+ \mu^-$



Possible new physics  
in charm FCNC


Observe:  
 $D^+ \rightarrow \varphi \pi^+$   
with  $\varphi \rightarrow \mu^+ \mu^-$



Put limits on:  
 $D^+ \rightarrow \pi^+ \mu^+ \mu^-$  with anti- $\varphi$  cut

$$\Rightarrow \text{BR}(D^+ \rightarrow \pi^+ \mu^+ \mu^-) < 4.7 \times 10^{-6}$$

Tevatron experiments make a lot of charm!



# $\psi(3770) \rightarrow hadrons$

## CLEO-c: Closing the gap

$$\sigma(e^+e^- \rightarrow D\bar{D}) = 6.39 \pm 0.10^{+0.17}_{-0.08} \text{ nb}$$

*PRL 95(2005)121801*

$$\sigma(e^+e^- \rightarrow hadrons) = 6.38 \pm 0.08^{+0.41}_{-0.30} \text{ nb}$$

*PRL 96(2006)092002*


Upper limit on gap is  $\approx 10\%$ . Other observed modes  $\approx 2\%$ .

## BES III: Resonance scan of $\psi(3770)$

*hep-ex/0605105 and hep-ex/0605107*



Find room for possible non- $DD$  contribution of  $\approx 16 \pm 8\%$ .

Consistent with CLEO-c, worth more study.







# $^1P_1$ Charmonium: The $h_c(3525)$



PRD 72(2005)032001

PRD 72(2005)092004




$$e^+e^- \rightarrow \psi(2S) \rightarrow \pi^0 h_c$$

$$h_c \rightarrow \gamma \eta_c$$




# Radial $\chi_{cJ}$ Excitations?



$$e^+e^- \rightarrow J/\psi + X(3940)$$

Consistent with  $J^{PC}=0^{++}$



$$\Upsilon\Upsilon \rightarrow Z(3930) \rightarrow D\bar{D}$$

Consistent with  $J^{PC}=2^{++}$


# The $Y(4260)$




# BABAR

PRL 95(2005)142001


$e^+e^- \rightarrow \pi^+\pi^- J/\psi$  using  
“radiative return”



New States



PRL 96(2006)  
162003






# BABAR

Preliminary

@ La Thuile 2006

Lattice QCD

# $D_s$ Decay Constant




Lattice QCD calculates:

$$f_{D_s} = 249 \pm 3 \pm 16 \text{ MeV}$$

*PRL 95(2005)122002*

Uses “charm tagging” to find leptonic decays of  $D_s$  from  $D_s^* \rightarrow \gamma D_s$



BaBar finds:



$$f_{D_s} = 279 \pm 17 \pm 6 \pm 19 \text{ MeV}$$


*Third error from branching ratio for  $D_s \rightarrow \phi \pi$ , recent from BaBar and which CLEO-c will measure to higher precision.*

# CLEO-c: D Tagging


Example:  $D \rightarrow \mu \nu$

$e^+ e^- \rightarrow D^+ D^-$





# D<sup>+</sup> Decay Constant



*Signal in Missing Mass*

CLEO-c finds:

$$f_{D^+} = 222.6 \pm 16.7^{+2.8}_{-3.4} \text{ MeV}$$

Lattice QCD calculates:

$$f_D = 201 \pm 3 \pm 17 \text{ MeV}$$

PRL 95(2005)122002

- Important test of actions that use “staggered fermions.”
- Same for determinations of  $f_{D_s}$ .
- More results to come!




# BABAR



# Semileptonic Form Factors

Belle: hep-ex/0604049



Lattice QCD

*Nucl.Phys.Proc.Suppl. 129(2004)334*

Pole model


ISGW2 prediction

*Low background results from  
CLEO-c are on the way.*




# Semileptonic Decay



Look for “Wrong Sign” lepton using  $D^{*\pm} \rightarrow D^0 \pi^\pm$  to tag flavor



“Right” Sign



“Wrong” Sign as a function of decay time



Upper limit on mixing rate  $r_D < 1.0 \times 10^{-3}$





# D<sup>0</sup> → Kπ DCS & Mixing



## Belle: Time Dependence

PRL 96(2006)151801


## CDF: Wrong sign Kπ

hep-ex/0605027

$$R_D = 0.405 \pm 0.021 \pm 0.011\%$$





$$R_D = 0.377 \pm 0.008 \pm 0.005\%$$




# Limits on Mixing Parameters

*D. Asner, Review in 2006 Particle Data Group compilation*



Experiment	Mode	A <sub>CP</sub> (%)	Notes
BaBar	$D^+ \rightarrow K^+ K^- \pi^+$	$1.4 \pm 1.0 \pm 0.8$	Exploits resonant substructure
BaBar	$D^+ \rightarrow \varphi \pi^+$	$0.2 \pm 1.5 \pm 1.6$	
BaBar	$D^+ \rightarrow K^{*0} K^+$	$0.9 \pm 1.7 \pm 0.7$	
CLEO II.V	$D^0 \rightarrow \pi^+ \pi^- \pi^0$	$1^{+9}_{-7} \pm 8$	Dalitz plot
CDF	$D^0 \rightarrow K^+ K^-$	$2.0 \pm 1.2 \pm 0.6$	Direct CP
CDF	$D^0 \rightarrow \pi^+ \pi^-$	$1.0 \pm 1.3 \pm 0.6$	
FOCUS	$D^0 \rightarrow K^+ K^- \pi^+ \pi^-$	$1.0 \pm 5.7 \pm 3.7$	Triple correlations to get at <i>T</i> -violation
FOCUS	$D^+ \rightarrow K^0 K^+ \pi^+ \pi^-$	$2.3 \pm 6.2 \pm 2.2$	
FOCUS	$D_s^+ \rightarrow K^0 K^+ \pi^+ \pi^-$	$-3.6 \pm 6.7 \pm 2.3$	




hep-ex/0606045

D<sup>0</sup> → K\* K:

A Tool for B<sup>±</sup> → D<sup>0</sup> K<sup>±</sup>

D<sup>0</sup> → K<sup>+</sup> K<sup>-</sup> π<sup>0</sup> Dalitz analysis



Destructive interference:

$$\delta_D = 332^\circ \pm 8^\circ \pm 11^\circ$$

Amplitude ratio:

$$\frac{|A(K^{*-} K^+)|}{|A(K^{*+} K^-)|} = 0.52 \pm 0.05 \pm 0.04$$

Submitted to Phys Rev D

# Quantum Correlations

For  $e^+e^- \rightarrow \bar{D}^0 D^0$  expect  $CP(\bar{D}^0 D^0) = -1$

This can be exploited in a number of ways, including extract CP content for multibody charm decays and searching for CP violation.

CLEO-c is studying the ways we can use this in our data, and looking forward to applying these ideas to new data samples.


BES III will be in an excellent position to capitalize!

Ref: D.Asner and W. Sun, *Phys.Rev.D*73(2006)034024

Statistical errors only!

D<sup>0</sup> Mixing and CP

# Preliminary



CP+

No QC

CP-

Data

	K <sup>+</sup> K <sup>-</sup>	π <sup>+</sup> π <sup>-</sup>	K <sub>S</sub> π <sup>0</sup> π <sup>0</sup>	K <sub>S</sub> π <sup>0</sup>
K <sup>+</sup> K <sup>-</sup>	5.2±0.4 -2.2±1.9	4.5±0.3 0.1±0.9	5.7±0.4 1.6±1.3	16.0±0.6 39.6±6.3
π <sup>+</sup> π <sup>-</sup>		1.1±0.2 0.2±1.4	2.2±0.2 1.6±1.3	5.8±0.4 14.0±3.7
K <sub>S</sub> π <sup>0</sup> π <sup>0</sup>			1.2±0.2 1.0±1.0	7.3±0.4 19.0±4.4
K <sub>S</sub> π <sup>0</sup>				9.7±0.5 3.0±1.7

Product CP+

Product CP-


# The Future

- Expect more from Belle, BaBar, CDF, and D0  
They produce lots of charm!
- CLEO-c will run through March 2008  
Expect  $\approx 3\text{M}$  D-pairs (charged and neutral)  
Also “thousands” of tagged  $D_s$  *Sneak Peek!*
- BES III coming on line in the next few years  
Data samples to be  $\approx 25\times$  CLEO-c
- Don't forget about LHCb, PANDA, ...

*Obrigado!*

*Also, thanks to all the experiments, and especially to R. Briere!*

# CLEO-c Preliminary: $D_s$ Hadronic Decays



Mode	CLEO-c (%)	PDG 2004 fit (%)
$\mathcal{B}(K_S K^+)$	$1.28^{+0.13}_{-0.12} \pm 0.07$	$1.8 \pm 0.55$
$\mathcal{B}(K^- K^+ \pi^+)$	$4.54^{+0.44}_{-0.42} \pm 0.25$	$4.3 \pm 1.2$
$\mathcal{B}(K^- K^+ \pi^+ \pi^0)$	$4.83^{+0.49}_{-0.47} \pm 0.46$	—
$\mathcal{B}(\pi^+ \pi^+ \pi^-)$	$1.02^{+0.11}_{-0.10} \pm 0.05$	$1.00 \pm 0.28$