

UNIVERSALITY & LFV IN UPSILON DECAYS AT CLEO

J.E. DUBOSCQ

CORNELL

JED@MAIL.LEPP.CORNELL.EDU

FOR THE CLEO COLLABORATION

TAU06 SEPT 19 2006

TOPICS

- CLEOIII DETECTOR AND DATA
- A SEARCH FOR THE LEPTON FLAVOR VIOLATING DECAY $\Upsilon \rightarrow \mu\tau$
- TEST OF LEPTON UNIVERSALITY IN $\Upsilon \rightarrow \tau\tau$ AND $\Upsilon \rightarrow \mu\mu$

THE CLEOIII DETECTOR

2230104-001

CLEOIII


EXCELLENT
CALORIMETER COVERAGE
AND RESOLUTION

EXCELLENT TRACKING
COVERAGE AND
RESOLUTION

RING IMAGING
CERENKOV & DE/DX
FOR PID

MUON CHAMBERS



+ A GREAT ACCELERATOR
TEAM CESR



CLEO UPSILON DATA

CLEOIII HAD THE LARGEST WORLD SAMPLE OF CLEAN Υ EVENTS BELOW B THRESHOLD (CF BELLE 3S)

ALSO HAS OFF-RESONANCE DATA + SCAN DATA + DATA AT Υ (5S)



SEARCH FOR LFV
VIOLATION IN $\Upsilon \rightarrow \mu\tau$

LEPTON FLAVOR VIOLATION


- WE DON'T ANNIHILATE WHEN WE SHAKE HANDS.
- BARYON ASYMMETRY GENERATED VIA SAKHAROV CONDITIONS: B & C & CP VIOLATION, + UNIVERSE OUT OF THERMAL EQUILIBRIUM FOR A WHILE
- B, L ARE ACCIDENTAL SYMMETRIES OF SM
- B-L IS HOWEVER CONSERVED
- MAYBE B IS VIOLATED BECAUSE L IS VIOLATED AND MAYBE THIS IS RELATED TO LFV ?

LFV FOR τ AND Υ DECAYS

■ τ DECAY LFV TALKS ABOUT Υ LFV

S.NUSSINOV, R.D.PECCEI, X.M. ZHANG PRD63(2000), 016003



$$B(\tau \rightarrow 3 \mu) < 10^{-6} \Rightarrow B(\Upsilon \rightarrow \mu \tau) < 10^{-2}$$



SUSY LOOPS


W.J. HUO, C.X. YUE, T.F. HENG, PRD67(2003), 114001

$$B(\Upsilon \rightarrow \mu \tau) < 2.2 \times 10^{-9}$$



LEPTOQUARK Z'


$$B(\Upsilon \rightarrow \mu \tau) < 1.3 \times 10^{-8}$$



LFV FOR τ , Υ DECAYS

Z. SILAGADZE, PHYS. SCRIPTA 64(2001),
128

GENERIC 4 FERMION
COUPLING α_N AT SCALE Λ
ADDED TO SM TO GET LFV




$$\frac{B(\Upsilon \rightarrow \mu\tau)}{B(\Upsilon \rightarrow \mu\mu)} \propto (\alpha_N/\alpha)^2 (M_\Upsilon/\Lambda)^4$$

LFV: THE ANALYSIS

- SEARCH FOR $\Upsilon \rightarrow \mu\tau$, $\tau \rightarrow e\nu\nu$
- 2 TRACKS: μ (MUON ID), e (E/P, DE/DX) μ NEAR EBEAM
- EXTENDED MAX LIKELIHOOD $\mathcal{L} = e^{-N_{evnt}} \Pi_{evnt} (\sum N_i \mathcal{P}_i(X|S))$
- USE PRODUCT PDF: $\mathcal{P}(p_\mu) \times \mathcal{P}(p_e) \times \mathcal{P}(dE/dx(e)) \times \mathcal{P}(E/p(e))$
- SUM OVER: SIGNAL LFV DECAY \oplus $\tau\tau$ \oplus $\mu\mu$ (γ) w/
HARD γ \oplus $\mu\mu$ WITH μ DECAY TO ELECTRON
- $\Upsilon(4S)$, OFF RES USED AS CALIBRATION & CONTROL SAMPLES

BACKGROUND & SIGNAL REGIONS



SIGNAL REGION


UPS(4S) DATA

$\mu\mu$ + HARD γ
 γ HITS CC AND
FAKES E/P

$\mu\mu$ & μ DECAY
IN FLIGHT
ABOUT 100 IN
 $\gamma(4S)$ DATA

EXTRACT MOST
PDFS FROM γ
(4S) DATA

FITS TO $\Upsilon(1S)$ DATA



- $\tau\tau$
- $\mu\mu$ HARD γ
- $\mu\mu$, DECAY IN FLIGHT

PRELIMINARY LFV RESULTS

RESONANCE	$\Upsilon(1S)$	$\Upsilon(2S)$	$\Upsilon(3S)$
EFFICIENCY	8.9%	8.9%	8.9%
NEVENTS	< 10.0	< 10.7	< 8.5
$B(\Upsilon \rightarrow \mu\tau)(10^{-6})$	< 6.2	< 25	< 22
$B(\Upsilon \rightarrow \mu\tau)/B(\Upsilon \rightarrow \mu\mu)$	< 0.023%	< 0.17%	< 0.13%

ALL LIMITS ARE 90% CL UL

- **LARGEST SYST: PDF SHAPES & CORRELATIONS**
- **FIRST LIMITS ON $\Upsilon \rightarrow \tau\mu$**
- **THESE BRs SET A LOWER LIMIT OF ≈ 1 TEV ON GENERIC LFV SCALE (ASSUMING STRONG COUPLING)**

UNIVERSALITY IN Υ
DECAYS WITH $\Upsilon \rightarrow \mu\mu, \tau\tau$


$\Upsilon \rightarrow \tau\tau$ MOTIVATION

- CLEO HAS MEASURED $B(\Upsilon(nS) \rightarrow \mu\mu)$, $\Gamma(ee \rightarrow \Upsilon(nS))$
 $N=1,2,3$ PRL94:012001(2005) HEP-EX/0409027
- $\Upsilon \rightarrow \tau\tau$ ROUNDS OUT THIS SERIES
- $B(\Upsilon(1S) \rightarrow \tau\tau)$ KNOWN TO $\approx 10\%$ IN PDG
- $\Upsilon(2S) \rightarrow \tau\tau$ “OBSERVED” BR=1.8+/-1.7%
- $\Upsilon(3S)$ NOT YET SEEN

$\Upsilon \rightarrow \tau\tau$ Motivation

- NAIVE UNIVERSALITY:

- $B(\Upsilon \rightarrow ee) = B(\Upsilon \rightarrow \mu\mu) = B(\Upsilon \rightarrow \tau\tau)$



- IF THIS AIN'T THE CASE, THERE'S SOME EXPLAINING TO DO

- SANCHIS-LOZANO: HIGGS SEARCHES HAVE A BLIND SPOT NEAR THE Υ CF NEXT TALK, HEP-PH/0307313

- THE DECAY CHAIN $\Upsilon \rightarrow \gamma \eta_b$, ($\eta_b \rightarrow A^0$), $A^0 \rightarrow \tau\tau$ COULD ALTER $N(\Upsilon \rightarrow \tau\tau)/N(\Upsilon \rightarrow \mu\mu)$, IF γ SOFT & UNDETECTED

GOAL: $B(\Upsilon \rightarrow \tau\tau)/B(\Upsilon \rightarrow \mu\mu)$

- ISOLATE $\mu\mu$ AND $\tau\tau$ SIGNALS AT & BELOW $\Upsilon(N S)$,
 $N=1..4$

- APPLY ON -S*OFF TO $\Upsilon(4S)$ DATA FOR $\mu\mu$ AND $\tau\tau$ -
EXPECT $B(4S \rightarrow \ell\ell) \approx 0$ $S = \mathcal{L}_{ON}/\mathcal{L}_{OFF} (E_{OFF}/E_{ON})^2$

- FIND OFF RESONANCE $\sigma(e^+e^- \rightarrow \tau\tau)$; COMPARE WITH
 $\sigma(e^+e^- \rightarrow \mu\mu)$

- APPLY ON-S*OFF TO 1S, 2S, 3S DATA FOR $\mu\mu$ and
 $\tau\tau$

- EXTRACT $R = N(\Upsilon \rightarrow \tau\tau)/N(\Upsilon \rightarrow \mu\mu)$

- GET BRANCHING RATIO $B(\Upsilon \rightarrow \tau\tau)$ USING $B(\Upsilon \rightarrow \mu\mu)$

TECHNIQUE CHECK

TECHNIQUE CHECK


HOW TO FIND $\Upsilon \rightarrow \tau\tau, \mu\mu$

- USE $\mu\mu$ CUTS SIMILAR TO PREVIOUS $\Upsilon \rightarrow \mu\mu$ STUDY
- USE 1 PRONG DECAYS : $B(\tau \rightarrow 1 \text{ PRONG}) \approx 75\%$
- 2 TRACK EVENTS, PASSING GENERIC $\tau\tau$ MISSING MOMENTUM, ENERGY CUTS (NEUTRINOS!)
- CLASSIFY TRACKS AS e, μ, X
- INCLUDE NEUTRAL ENERGY, SHOWER CUTS


COSMICS ARE REJECTED

ON & S*OFF AT THE $\Upsilon(4S)$


$\Upsilon(4S) E(\tau\tau)/E_{CM}$ | vs | modes




$\Upsilon(4S) E(\tau\tau)/E_{CM}$ | vs X modes




$\Upsilon(4S) E(\tau\tau)/E_{CM}$ other modes



$\Upsilon(4S) E(\tau\tau)/E_{CM}$ All modes



MuPair Mass On, Off Res



ON-S*OFF WORKS AT $\Upsilon(4S)$

NO EVIDENCE FOR NON- $1/E^2$ BACKGROUND

OFF RES $\sigma(\tau\tau)/\sigma(\mu\mu)/\text{EXP}$


BREAKDOWN BY τ DECAY CHANNELS

$$\frac{\sigma^{\text{theory}}(e^+e^- \rightarrow \tau\tau)}{\sigma^{\text{theory}}(e^+e^- \rightarrow \mu\mu)} = 0.82$$

$(X,Y) = P(X) > P(Y)$

AVERAGE OVER
LEPTON MODES

OVERALL AVERAGE






ALL DECAY CHANNELS AGREE




WE CAN RECONSTRUCT $ee \rightarrow \tau\tau, \mu\mu$

ON AND S*OFF RES FOR $\tau\tau$, $\mu\mu$

TOTAL $\tau\tau$ RECONSTRUCTED ENERGY / E_{CM} :



MASS OF $\mu\mu$ / E_{CM} :





DATA REMAINING AFTER ON-S*OFF SUBTRACTION SHOULD BE ALL DUE TO Υ DECAYS


GETTING $N(\Upsilon \rightarrow \ell\bar{\ell})$

- DATA AFTER ON-S*OFF IS A SUM OF SIGNAL $\Upsilon \rightarrow \ell\bar{\ell}$, CASCADE TO LOWER $\Upsilon \rightarrow \ell\bar{\ell}$, OTHER Υ DECAYS.
- MEASURE $\text{BR}(\Upsilon(1S) \rightarrow \ell\bar{\ell})$ TO SCALE MC CASCADE BGD FOR $\Upsilon(2S)$ DECAYS, ITERATE FOR $\Upsilon(3S)$.
- USE KORALB FOR $\Upsilon \rightarrow \tau\bar{\tau}$ WITH ISR TURNED OFF
 - Υ HAS THE QUANTUM NUMBERS OF THE PHOTON
 - KORALB HAS HELICITY CORRELATIONS

A SPRINKLING OF PLOTS




**TOTAL ENERGY
DISTRIBUTION FOR 3
DIFFERENT MODES
AT 3 RESONANCES
FOR $\tau\tau$, ASSUMING
UNIVERSALITY**




NEED MORE CONVINCING ?


Ptag Ups1s Tau all




Psig Ups1s Tau all



Ecc Ups1s Tau all



Nshw Ups1s Tau all



GOOD AGREEMENT W/ MC ACROSS RESONANCES, FINAL STATES, AND KINEMATIC QUANTITIES

HOW MANY EVENTS ?

SUM OF ALL τ DECAY MODES

	1S	2S	3S
On-S*Off	61697 \pm 1536	25085 \pm 1399	16290 \pm 1522
background	1556 \pm 83	3334 \pm 593	1536 \pm 474
$\epsilon(\tau\tau)$	11.2 \pm 0.1%	11.3 \pm 0.1%	11.1 \pm 0.1%
$N(\tau\tau)/\epsilon (10^3)$	537 \pm 14	193 \pm 12	132 \pm 13
$N(\mu\mu)/\epsilon (10^3)$	527 \pm 15	185 \pm 11	126 \pm 11

STAT, MC STAT ERRORS INC

$$\mathcal{R} = B(\Upsilon \rightarrow \tau\tau) / B(\Upsilon \rightarrow \mu\mu)$$

FINAL
RESULT

$$\mathcal{R}(1S) = 1.02 \pm 0.02 \pm 0.05$$

$$\mathcal{R}(2S) = 1.04 \pm 0.04 \pm 0.05$$

$$\mathcal{R}(3S) = 1.07 \pm 0.08 \pm 0.05$$


LARGEST SYST FROM τ SELECTION CRITERIA (2.9%) AND TRIGGER (1.6%)

ALMOST ALL STAT ERROR FROM ON/OFF SUBTRACTION

SUBMITTED TO PRL
HEP-EX/0607019

\mathcal{R} by Decay Mode

$$\mathcal{R} = B(\Upsilon \rightarrow \tau\tau) / B(\Upsilon \rightarrow \mu\mu)$$



EXTRACTING $B(\Upsilon \rightarrow \tau\tau)$

- USE CLEO'S PUBLISHED $B(\Upsilon \rightarrow \mu\mu)$ - CF PRL94,012001(2005)
- AVOID SYST ERROR DOUBLE COUNTING

$$B(\Upsilon(1S) \rightarrow \tau\tau) = 2.54 \pm 0.04 \pm 0.12 \%$$

τ stat *SYST+ μ STAT*

$$B(\Upsilon(2S) \rightarrow \tau\tau) = 2.11 \pm 0.07 \pm 0.13 \%$$


$$B(\Upsilon(3S) \rightarrow \tau\tau) = 2.55 \pm 0.19 \pm 0.15 \%$$

WHAT ABOUT η_B /HIGGS?

- \mathcal{R} IS CONSISTENT WITH 1 - NO BIG SIGNAL
- ATTRIBUTE DEVIATION $\mathcal{R}(1S)-1$ TO “HIGGS”

$$B(\Upsilon \rightarrow \gamma \eta_B, \eta_B \rightarrow A^0, A^0 \rightarrow \tau\tau) < 0.27\% \quad 90\% \text{ CL UL}$$

- LOOK AT Υ SPECTRUM
 - NO OBVIOUS SPIKE IN THE 1S SPECTRUM
 - EXPECTED REGION IS FALLING FAST - HARD SYST
 - NO OBVIOUS SPIKE IN 2S, 3S SPECTRA



CONCLUSIONS

CLEO HAS:

- SEARCHED FOR LFV GIVES $BR(\Upsilon \rightarrow \mu\tau) \approx < 10^{-5}$ 90%CL UL
- MEASURED $B(\Upsilon \rightarrow \tau\tau)/B(\Upsilon \rightarrow \mu\mu)$ CONSISTENT WITH 1
- MEASURED $B(\Upsilon \rightarrow \tau\tau)$
 - CONSISTENT WITH PDG AT 1S (BUT LOWER)
 - BEST VALUE FOR 2S
 - FIRST VALUE FOR 3S
- SET LIMIT ON CP ODD HIGGS IN $\Upsilon(1S)$ REGION