



Search for $B_s \rightarrow \mu^+ \mu^-$ Decays



Walter Hopkins

Cornell University

Lattice Meets Experiment 2010



Motivation

- $B_s \rightarrow \mu^+ \mu^-$ can only occur through higher order FCNC diagrams in Standard Model (SM)
- This decay is not only suppressed by the GIM Mechanism but also by helicity
- SM predicts very low rate with little SM background ($\mathcal{BR}(B_s \rightarrow \mu^+ \mu^-) = (3.86 \pm 0.57) \times 10^{-9}$, M. Artuso et al, Eur. Phys. J. C57)
- Super symmetry (SUSY) models predict enhancement of this decay by $\tan\beta^6$
- Clean experimental signature $\rightarrow \tau$'s would have stronger coupling but experimentally difficult




Detector

- Reporting on 3.7 fb^{-1} CDF result, first shown in Fall 2009
- Secondary vertex ID with excellent Silicon tracker: $\sigma_{p_t}/p_t^2 \sim 0.15\%$ and $\sigma_{vtx} = 30 \mu\text{m}$
- Muon System



Experimental Challenges

- Large background at hadron collider
 - Must reduce large background around dimuon mass of $m_{B_s} = 5.37$ GeV
 - Analysis requirements: Design an effective discriminant, determine the efficiency for signal, and estimating the background level



Central-Central (CMU) and Central-Forward (CMX) Di-muon Trigger

- **Central:** $p_T > 2.0$ GeV and $|\eta| < 0.6$ – **Forward:** $p_T > 2.2$ GeV and $0.6 < |\eta| < 1.0$
- p_T cuts restrict us to well understood trigger regions



Basic Quality Cuts

- Tracker tracks with hits in 3 silicon layers
- Likelihood and dE/dx based muon Id
- Vertex Quality
- Loose preselection and analysis cuts
 - $p_T(\mu^+\mu^-) > 4.0$ GeV; 3D Decay length significance > 2
 - Loose Isolation and opening angle (pointing) cuts

Still background dominated after a reduction of events of 4 orders of magnitude

Analysis Method

- Measure rate of $B_s \rightarrow \mu^+ \mu^-$ relative to $B^+ \rightarrow J/\psi K^+$, $J/\psi \rightarrow \mu^+ \mu^-$
- Apply same selection to find $B^+ \rightarrow J/\psi K^+$
- Systematic uncertainties will cancel in ratio \Rightarrow e.g. dimuon trigger efficiency is the same for both modes
- D0 total B^+ yield: 5728 ± 85 (with 5 fb^{-1})





$$BR(B_s \rightarrow \mu^+ \mu^-) = \frac{N_{B_s} \epsilon_{B^+}^{trig} \epsilon_{B^+}^{reco}}{N_{B^+} \epsilon_{B_s}^{trig} \epsilon_{B_s}^{reco}} \cdot \frac{\alpha_{B^+}}{\alpha_{B_s}} \frac{1}{\epsilon_{NN}^{B^+}} \cdot \frac{f_{J/\psi}}{f_s} \cdot BR(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+)$$

From Data, From MC, From PDG

- Estimate acceptances and efficiencies
- Identify variables that discriminate signal and background
- Make multivariate discriminant, for background rejection
 - Optimized with Pythia signal MC and data mass sideband
 - Validate in B^+ sample
- Estimate Background
 - Combinatoric background
 - Peaking background: $B \rightarrow hh$



Discriminating Variables (CDF)

- Invariant mass of muons with 2.5σ window, $\sigma=24$ MeV
- 3 Secondary vertex related variables
 - $\lambda = c\tau$, proper decay time
 - $\frac{\lambda}{\sigma_\lambda}$
 - $\Delta\alpha = |\phi_B - \phi_{vtx}|$
- Isolation: $\frac{p_T(B)}{\Sigma p_T(trks) + p_T(B)}$
- Transverse momentum of B and lower momentum muon



Discriminating Variables: Neural Network (CDF)

- Combined all variables except mass in neural network
- Unbiased optimization based on MC signal and sideband data
- Extensively tested for mass bias



Control Regions (CDF)

- Test background estimates in blinded signal region with independent data samples
- Compare predicted vs. observer background events for multiple NN events

Regions

- OS-: Opposite sign muons with negative proper decay length
- SS+ and SS-: Same sign muons, positive and negative decay length
- FM: OS- & OS+ with one μ failing muon id and loose vertex cuts

sample	NN cut	CMU-CMU			CMU-CMX		
		pred	obsv	prob(%)	pred	obsv	prob(%)
OS-	$0.80 < \nu_{NN} < 0.95$	$275 \pm (9)$	287	26	$310 \pm (10)$	304	39
	$0.95 < \nu_{NN} < 0.995$	$122 \pm (6)$	121	46	$124 \pm (6)$	148	3.2
	$0.995 < \nu_{NN} < 1.0$	$44 \pm (4)$	41	36	<u>$31 \pm (3)$</u>	<u>50</u>	<u>0.4</u>
SS+	$0.80 < \nu_{NN} < 0.95$	$2.7 \pm (0.9)$	1	29	$2.7 \pm (0.9)$	0	10
	$0.95 < \nu_{NN} < 0.995$	$1.2 \pm (0.6)$	0	34	$1.2 \pm (0.6)$	1	66
	$0.995 < \nu_{NN} < 1.0$	$0.6 \pm (0.4)$	0	55	$0.0 \pm (0.0)$	0	-
SS-	$0.80 < \nu_{NN} < 0.95$	$8.7 \pm (1.6)$	9	49	$5.7 \pm (1.6)$	2	11
	$0.95 < \nu_{NN} < 0.995$	$3.0 \pm (1.0)$	4	36	$3.6 \pm (1.0)$	2	34
	$0.995 < \nu_{NN} < 1.0$	$0.9 \pm (0.5)$	0	43	$0.3 \pm (0.3)$	0	70
FM+	$0.80 < \nu_{NN} < 0.95$	<u>$169 \pm (7)$</u>	<u>169</u>	<u>50</u>	$73 \pm (5)$	64	19
	$0.95 < \nu_{NN} < 0.995$	$55 \pm (4)$	43	9	$19 \pm (2)$	18	49
	$0.995 < \nu_{NN} < 1.0$	$20 \pm (2)$	20	48	$3.6 \pm (1.0)$	3	53

Expected Sensitivities (CDF)

- Single event sensitivity is at SM level ($= 3.86 \times 10^{-9}$)
- Largest uncertainty from $\frac{f_u}{f_s}$





	CMU-CMU		CMU-CMX	
$(\alpha_{B^+}/\alpha_{B_s})$	0.300 ± 0.018	(±6%)	0.196 ± 0.0014	(±7%)
$(\epsilon_{B^+}^{trig}/\epsilon_{B_s}^{trig})$	0.99935 ± 0.00012	(-)	0.97974 ± 0.00016	(-)
$(\epsilon_{B^+}^{reco}/\epsilon_{B_s}^{reco})$	0.82 ± 0.03	(±4%)	0.83 ± 0.03	(±4%)
$\epsilon_{B_s}^{NN} (NN > 0.80)$	0.776 ± 0.047	(±6%)	0.789 ± 0.047	(±6%)
N_{B^+}	14300 ± 170	(±1%)	5460 ± 110	(±2%)
f_u/f_s	3.86 ± 0.59	(±15%)	3.86 ± 0.59	(±15%)
$BR(B^+ \rightarrow J/\psi K^+ \rightarrow \mu^+ \mu^- K^+)$	$(5.94 \pm 0.21) \times 10^{-5}$	(±4%)	$(5.94 \pm 0.21) \times 10^{-5}$	(±4%)
SES (All bins)	5.1×10^{-9}	(±18%)	8.5×10^{-9}	(±19%)
SES (Combined)	3.2×10^{-9} (±18%)			

Neural Network

- 3 NN bins, majority of sensitivity comes from highest bin
- Treated separately → Different Signal/Background
- Lower NN bins added → 50% increase in efficiency and improved sensitivity
- Expected Signal: $NN > 0.8 \rightarrow 1.2$ events




Background

- Combinatoric Background
 - Estimated with linear fit to sideband
 - Use p0 and exp fit in highest NN bin for syst. error estimation
- $B \rightarrow hh$
 - Peaks in signal region
 - Use $B_{S(d)} \rightarrow hh$ MC to estimate acceptance and convolute with muon fake rate from data using D^* tagged to $D \rightarrow K\pi$
 - Order of magnitude larger for B_d vs. B_s
 - For $NN > 0.995$ in B_d mass window 0.81 events




Dimuon Mass vs NN

CDF at 3.7 fb^{-1}




D0 at 2 fb^{-1}




- Systematic uncertainties included
- CDF has worlds best limit at 4.3×10^{-8} @ 95% CL with 3.7 fb^{-1}
- D0 expected sensitivity with 5 fb^{-1} : 5.3×10^{-8} @ 95% CL
- Last published D0 limit with 2 fb^{-1} : 9.3×10^{-8} @ 95% CL

95% CL Limits on $\mathcal{B}(B_s \rightarrow \mu\mu)$



	$\mathcal{B}(B_s \rightarrow \mu^+ \mu^-)$ 90%	95%	$\mathcal{B}(B_d \rightarrow \mu^+ \mu^-)$ 90%	95%
Expected \mathcal{B}	2.7×10^{-8}	3.3×10^{-8}	7.2×10^{-9}	9.1×10^{-9}
Observed \mathcal{B}	3.6×10^{-8}	4.3×10^{-8}	6.0×10^{-9}	7.6×10^{-9}



CDF BR($B_s \rightarrow \mu^+ \mu^-$) Projection



CDF working to update and improve analysis for later this year

- More data, up to 6.7 fb^{-1}
- Apply improved dE/dx calibration
- Increase acceptance by introducing more detector regions, now better understood

Future: LHCb



- B Focused forward experiment
→ many boosted B's
- Will reach SM limits quickly
with less luminosity
- Similar discriminating variables

Plot shows for $E_{cm}=14$ TeV

Conclusion

CDF Preliminary Results with 3.7 fb^{-1}

$$\mathcal{BR}(B_s \rightarrow \mu^+ \mu^-) = 4.3 \times 10^{-8} \text{ at 95\% CL}$$

$$\mathcal{BR}(B_d \rightarrow \mu^+ \mu^-) = 7.6 \times 10^{-9} \text{ at 95\% CL}$$

- Reached sensitivity at the 3.2×10^{-9} level
- Set the world's best limits for both B_s and B_d in these modes
- Probing new parameter space across a variety of New Physics models
- D0 updating their analysis with 5 fb^{-1}
- LHCb projects Tevatron limit with 0.15 fb^{-1} at $E_{cm} = 14 \text{ TeV}$

mSUGRA at $\tan\beta = 50$
Arnowitz, Dutta, et al., PLB 538 (2002) 121

